

Epigenetics

ISSN: 1559-2294 (Print) 1559-2308 (Online) Journal homepage: https://www.tandfonline.com/loi/kepi20

Differential DNA methylation and $PM_{2.5}$ species in a 450K epigenome-wide association study

Lingzhen Dai, Amar Mehta, Irina Mordukhovich, Allan C. Just, Jincheng Shen, Lifang Hou, Petros Koutrakis, David Sparrow, Pantel S. Vokonas, Andrea A. Baccarelli & Joel D. Schwartz

To cite this article: Lingzhen Dai, Amar Mehta, Irina Mordukhovich, Allan C. Just, Jincheng Shen, Lifang Hou, Petros Koutrakis, David Sparrow, Pantel S. Vokonas, Andrea A. Baccarelli & Joel D. Schwartz (2017) Differential DNA methylation and $PM_{2.5}$ species in a 450K epigenome-wide association study, Epigenetics, 12:2, 139-148, DOI: 10.1080/15592294.2016.1271853

To link to this article: https://doi.org/10.1080/15592294.2016.1271853

View supplementary material

Published online: 31 Jan 2017.

Submit your article to this journal

Article views: 1395

View related articles

View Crossmark data

Citing articles: 18 View citing articles

Full Terms & Conditions of access and use can be found at
https://www.tandfonline.com/action/journalInformation?journalCode=kepi20

EPIGENETICS
2017, VOL. 12, NO. 2, 139–148
http://dx.doi.org/10.1080/15592294.2016.1271853

Taylor & Francis
Taylor & Francis Group

RESEARCH PAPER

Differential DNA methylation and $PM_{2.5}$ species in a 450K epigenome-wide association study

Lingzhen Dai[a], Amar Mehta[a], Irina Mordukhovich[a], Allan C. Just [b], Jincheng Shen[c], Lifang Hou[d], Petros Koutrakis[a], David Sparrow[e], Pantel S. Vokonas[e], Andrea A. Baccarelli [a], and Joel D. Schwartz[a]

[a]Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA, USA; [b]Department of Preventive Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA; [c]Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, USA; [d]Department of Preventive Medicine, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA; [e]Veterans Affairs Normative Aging Study, Veterans Affairs Boston Healthcare System, Department of Medicine, Boston University School of Medicine, Boston, MA, USA

ABSTRACT

Although there is growing evidence that exposure to ambient particulate matter is associated with global DNA methylation and gene-specific methylation, little is known regarding epigenome-wide changes in DNA methylation in relation to particles and, especially, particle components. Using the Illumina Infinium HumanMethylation450 BeadChip, we examined the relationship between one-year moving averages of $PM_{2.5}$ species (Al, Ca, Cu, Fe, K, Na, Ni, S, Si, V, and Zn) and DNA methylation at 484,613 CpG probes in a longitudinal cohort that included 646 subjects. Bonferroni correction was applied to adjust for multiple comparisons. Bioinformatics analysis of the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment was also performed. We observed 20 Bonferroni significant (P-value $< 9.4\times 10^{-9}$) CpGs for Fe, 8 for Ni, and 1 for V. Particularly, methylation at *Schlafen Family Member 11* (*SLFN11*) cg10911913 was positively associated with measured levels of all 3 species. The *SLFN11* gene codes for an interferon-induced protein that inhibits retroviruses and sensitizes cancer cells to DNA-damaging agents. Bioinformatics analysis suggests that gene targets may be relevant to pathways including cancers, signal transduction, and cell growth and death. Ours is the first study to examine the epigenome-wide association between ambient particles species and DNA methylation. We found that long-term exposures to specific components of ambient particle pollution, especially particles emitted during oil combustion, were associated with methylation changes in genes relevant to immune responses. Our findings provide insight into potential biologic mechanisms on an epigenetic level.

ARTICLE HISTORY
Received 8 June 2016
Revised 26 November 2016
Accepted 7 December 2016

KEYWORDS
Ambient particulate matter; DNA methylation; epigenome-wide association study; metals; pathway enrichment

Introduction

The associations between fine particulate matter (particles with an aerodynamic size of $\leq 2.5\ \mu$m, $PM_{2.5}$) and health outcomes are well established. It is estimated that $PM_{2.5}$ leads to 3.15 million premature deaths per year on a global scale.[1] Nationwide and multiple-community studies report that exposure to $PM_{2.5}$ is linked with an increased risk of hospital admissions and mortality in the US, and that certain $PM_{2.5}$ species may be more toxic than others.[2–5]

DNA methylation is an epigenetic process in which a methyl group is added to deoxycytosine bases to form deoxymethylcytosine. The literature to date indicates that DNA methylation may be an important pathway linking particulate matter (PM) to health outcomes.[6–10] For example, changes in patterns of DNA methylation may be associated with processes leading to cardiovascular diseases..[11,12] There is growing evidence that exposure to PM is associated with global DNA methylation and gene-specific methylation,[6,7,13,14] especially for exposure to metal-rich PM in occupational settings.[15,16] Yet, little is known regarding how PM exposure affects DNA methylation on a genome-wide level.

Measurements of DNA methylation at cytosine-guanine dinucleotide (CpG) loci have recently become available; for instance, the Illumina Infinium HumanMethylation450 BeadChip (450K) measures methylation at more than 450,000 CpG sites. This makes an epigenome-wide association study of the relationship between particles and DNA methylation possible. Several studies have investigated the epigenome-wide association between smoking, an important PM source, and DNA methylation. Using the 450K platform and Bonferroni correction, Joubert et al.[17] identified 26 CpGs with methylation changes in newborns related to maternal smoking during pregnancy in a large Norwegian birth cohort. Specifically, the authors documented changes at CpGs in *cytochrome p450 1A1 (CYP1A1)* and *aryl-hydrocarbon receptor repressor (AHRR)*, genes known to play a key role in the aryl hydrocarbon receptor (AhR) signaling pathway that mediates the detoxification of components of tobacco smoke. In a more recent study, the authors identified numerous loci associated with maternal smoking during pregnancy in a genome-wide consortium meta-analysis and found substantial persistence of such effects

CONTACT Lingzhen Dai ldai@mail.harvard.edu Department of Environmental Health, Harvard T.H. Chan School of Public Health, Landmark Center 4th floor, 401 Park Drive, Boston, MA 02215, USA.
Supplemental data for this article can be accessed on the publisher's website.
© 2017 Taylor & Francis Group, LLC

in later childhood.[18] Zeilinger et al.[19] conducted an epigenome-wide study comparing methylation levels among current, former, and never smokers in a population-based panel, and found widespread differences in the degree of site-specific methylation as a function of tobacco smoking. Moreover, they observed that the most significant associations with smoking were for DNA methylation sites in *AHRR*.

To date, very few epigenome-wide association studies have focused on exposures to ambient particles, with the exception of one recent study that identified several CpG sites associated with cumulative exposure to ambient particles among populations from Germany and the US.[20] To our knowledge, no epigenome-wide association studies have investigated the toxicity of specific PM species. Such research would be of great importance because epigenome-wide association research based on DNA methylation microarrays could provide insight into epigenetic mechanisms underlying associations between air particles and health outcomes and thereby improve our understanding of human disease. To our knowledge, our study is the first to examine epigenome-wide associations between $PM_{2.5}$ species and DNA methylation using data from a large longitudinal cohort.



Figure 1. Flowchart of study participants.

Methods

Study population

Study subjects were participants of the Normative Aging Study (NAS), a closed cohort that was established in 1963 by the Department of Veterans Affairs. This ongoing longitudinal study of aging enrolled 2,280 community-dwelling, healthy men living in the Greater Boston area.[21] Participants were free of known chronic medical conditions at enrollment and have undergone examinations every 3 to 5 y on a continuous rolling basis. All participants provided written informed consent. The study was reviewed and approved by the Institutional Review Boards of all participating institutions.

We restricted our analysis to study visits with DNA samples, which were collected from subjects starting in 1999. A comparison of visits with DNA samples and visits without showed no substantial differences with regard to body mass index, education, alcohol consumption, and smoking, although older participants were less likely to provide DNA samples (mean age 74.5 vs. 76.3 years; *P*-value of t-test in mean age < 0.05). In addition, since Caucasians comprise most of our study population (approximately 97%), we excluded non-Caucasians to reduce study heterogeneity that may be introduced by diverse genetic ancestry. We also excluded observations that did not have complete information on the exposures or covariates of interest. In the final analyses, samples from 646 subjects with 1,031 total visits were included. Of the 646 subjects, 314 presented for one visit, 279 presented for 2 visits, and 53 presented for 3 visits. The flowchart of our study participants is shown in Fig. 1.

DNA methylation

DNA samples were collected between 1999 and 2013. We used the QIAamp DNA Blood Kit (Qiagen, CA, USA) to extract DNA from buffy coat, and performed bisulfite conversion with the EZ-96 DNA Methylation Kit (Zymo Research, CA, USA). To minimize batch effects, we randomized chips across plates and randomized samples based on a 2-stage age stratified algorithm so that age distributed similarly across chips and plates.

We measured DNA methylation of CpG probes using Illumina Infinium HumanMethylation450 BeadChip.[22] Quality control analysis was performed, where samples with >1% of probes with a detection *P*-value > 0.05 were removed, and probes with *P*-value $> 5\%$ samples with a bead count < 3 or with >1% samples with a detection *P*-value > 0.05 were removed. The remaining samples were preprocessed using the Illumina-type background correction[23] and normalized with dye-bias[24] and BMIQ3 adjustments,[25] which were used to generate methylation values. Extreme outliers, defined by Tukey's method (i.e., $<25^{th}$ percentile $- 3 \times$ IQR or $> 75^{th}$ percentile $+ 3 \times$ IQR), were trimmed. As a result, 484,613 CpG probes were included in the working set.

We used the β value to indicate methylation level at each CpG site. The β value represents the ratio of the fluorescence of the methylated signal to the combined methylated and unmethylated signals; that is, $\beta =$ intensity of the methylated signal (M) / [intensity of the unmethylated signal (U) + intensity of the methylated signal (M) + 100].[22]

The methylation data has been uploaded to the database of Genotypes and Phenotypes (dbGaP; available at https://www.ncbi.nlm.nih.gov/gap) established by the National Institutes of Health, along with some other basic information (dbGaP accession phs000853.v1.p1).

Air pollution

We measured daily concentrations of $PM_{2.5}$ mass and species at the stationary monitoring supersite at the Harvard University Countway Library, Boston, MA, USA. $PM_{2.5}$ was assessed by

the Tapered Element Oscillating Microbalance (TEOM 1400a, Rupprecht & Patashnick Co.), and PM species were assessed using the Energy Dispersive X-ray Fluorescence Spectrometer (Epsilon 5, PANalytical).

To estimate long-term effects of air pollution on DNA methylation, we calculated one-year moving average concentrations for $PM_{2.5}$ and the following 11 species: Al, Ca, Cu, Fe, K, Na, Ni, S, Si, V, and Zn. These species are representative of different sources of particles: Al, Ca, Fe, and Si are major components of road dust; Cu and Zn originate from traffic emissions; Ni and V are tracers of oil combustion; S is a regional pollutant that is associated with coal combustion and to a lesser extent with metallurgic activities; Na is a major component of sea salt particles; K originates from wood burning and soil.

Statistical analysis

For each specific CpG site, we applied a linear mixed-effects model as described below,

$$E(Y_{ij}) = \beta_0 + \beta_1 * Spec_{ij} + \beta_2 * PM_{2.5ij} + \beta_3 * Age_{ij} + \cdots + \mu_i, \quad (1)$$

where Y_{ij} is the methylation level of subject i at visit j, $Spec_{ij}$ and $PM_{2.5ij}$ are one-year moving average species and $PM_{2.5}$ concentrations before visit j, Age_{ij} is subject i's age at visit j (this notation is relevant to all other covariates, since we used values assessed at each visit in our analyses), and μ_i is the random intercept that accounts for correlation within subject. The full formula can be found in the Supplemental Document.

In the analysis, we controlled for the following covariates *a priori* based on the relevant literature:[6,19,26,27] total $PM_{2.5}$ mass, cell proportions ($CD4^+$ T lymphocytes, $CD8^+$ T lymphocytes, natural killer cells, B cells, monocytes) estimated by the Houseman et al. method,[28–30] age, body mass index [BMI, computed as weight (in kilograms) divided by height (in square meters)], cigarette pack years, smoking status (never, ever), alcohol consumption ($\geq$ 2 drinks/day, $<$ 2 drinks/day), years of education, year and season (spring: March–May, summer: June–August, fall: September–November, winter: December–February). Potential batch effects were also considered, including plate, the position of the chip on plate, and the row and column position on the chip.

Since overall type I error inflates as the number of statistical tests increases, we used Bonferroni correction to adjust for multiple comparisons. This method, which accounts for the number of CpG sites and exposures tested, has been widely used in similar studies.[31–33] As a result, the adjusted level of significance in the current study was P-value $= 9.4\times 10^{-9}$, i.e., 0.05 / (484,613$\times$ 11). After identifying significant CpGs, we further checked whether they were non-unimodal in the dip test,[34] were cross-reactive,[35] or were located within 10 bases of SNPs relevant to European populations.[36]

Our study population was a heavily smoking population. Since we controlled for smoking status and pack years in the analysis, we believe there was sufficient adjustment, but we understand that the presence of a large number of smokers may obscure or distort associations. To address this, we have performed subset analyses among nonsmokers and smokers.

All statistical analyses were performed using R 3.1.2.[37]

Bioinformatics analysis

We performed Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis[38] using the genes targets of CpGs, identified by the Illumina annotation assignment, with Benjamini-Hochberg false discovery rates (FDR)[39] $<$1% and $<$5%, respectively. Bioinformatics analysis was performed using Database for Annotation, Visualization and Integrated Discovery (DAVID) Bioinformatics Resources 6.7.[31,32] Briefly, DAVID consists of an integrated biologic knowledgebase with wide annotation content coverage. It allowed us to identify the most relevant pathways associated with the gene lists we provided.

Results

Characteristics of study subjects are presented in Table 1. At baseline, the mean age was 74 y [standard deviation (SD) = 7]. The study population was well educated, with a mean of 15 y of education (SD = 3). Furthermore, 186 of the 646 (28.8%) individuals never smoked, and 131 (20.3%) had 2 or more drinks per day.

Table 2 shows the descriptive summary of concentrations of $PM_{2.5}$ mass and species. During the study period 1,999 – 2,013, the annual average $PM_{2.5}$ concentration was 10.5 (SD = 1.1) $\mu g/m^3$. Among the examined species, S accounted for the largest mass proportion (10%). The oil combustion tracers, Ni and V, accounted for 0.03% and 0.04% of $PM_{2.5}$ mass with mean concentrations of 0.0037 (SD = 0.0013) $\mu g/m^3$ and 0.0041 (SD = 0.0012) $\mu g/m^3$, respectively. We report a strong correlation between Ni and V (r = 0.96).

Table 3 presents CpGs that reached Bonferroni-corrected significance (P-value $< 9.4 \times 10^{-9}$) in relation to $PM_{2.5}$ species. After excluding CpGs that were non-unimodal in the dip test, were cross-reactive, or contained SNPs within 10 bases, we observed 20 significant CpGs for Fe, 8 for Ni, and 1 for V. Particularly, *Schlafen Family Member 11 (SLFN11)* cg10911913 was significant for all 3 metals. The ratio of methylation at this CpG site is expected to increase by 0.073, 0.031, and 0.044 per IQR increase in the one-year

Table 1. Characteristics of the study population.

Variable	First visit (n = 646)	All visits (n = 1,031)
Mean ± SD		
Age (years)	73.7 ± 6.8	74.7 ± 6.8
BMI (kg/m^2)	28.0 ± 4.1	28.0 ± 4.2
Education (years)	15.1 ± 3.0	15.2 ± 3.0
Cigarette smoking (pack years)	21.3 ± 25.5	21.3 ± 25.2
Number (%)		
Smoking status		
Never	186 (28.8)	298 (28.9)
Ever	460 (71.2)	733 (71.1)
Alcohol consumption (drinks/day)		
$\geq$ 2	131 (20.3)	200 (19.4)
$<$ 2	515 (79.7)	831 (80.6)

Table 2. Summary of one-year moving averages of $PM_{2.5}$ mass and species across all study visits (year 1999–2013).

			Correlation coefficient											
Pollutant	Mean ± SD (μg/m^3)	IQR (μg/m^3)	$PM_{2.5}$	S	Na	Al	Ca	Fe	Si	K	Zn	Cu	Ni	V
$PM_{2.5}$	10.5 ± 1.1	1.9	1	0.83	0.45	0.67	0.18	0.66	0.57	0.50	0.61	0.57	0.63	0.50
S	1.11 ± 0.13	0.08		1	0.81	0.57	0.17	0.55	0.41	0.32	0.51	0.54	0.69	0.63
Na	0.20 ± 0.01	0.01			1	0.37	0.11	0.24	0.19	0.18	0.20	0.39	0.42	0.45
Al	0.051 ± 0.008	0.009				1	0.60	0.89	0.96	0.65	0.51	0.87	0.80	0.71
Ca	0.032 ± 0.005	0.007					1	0.77	0.73	0.20	0.73	0.73	0.70	0.78
Fe	0.069 ± 0.010	0.012						1	0.91	0.51	0.76	0.88	0.93	0.86
Si	0.075 ± 0.016	0.022							1	0.61	0.57	0.85	0.79	0.72
K	0.041 ± 0.002	0.003								1	0.18	0.54	0.44	0.37
Zn	0.013 ± 0.003	0.004									1	0.64	0.72	0.73
Cu	0.0035 ± 0.0003	0.0004										1	0.80	0.78
Ni	0.0037 ± 0.0013	0.0011											1	0.96
V	0.0041 ± 0.0012	0.0015												1

moving averages of Fe, Ni, and V, respectively. We found strong seasonal patterns at this site for the 3 metals (Figure S1 in the Supplemental Material), with the highest methylation levels in the spring. We also found that methylation levels at *CASZ1* cg16238819, *CUEDC2* cg06024834, *ONECUT1* cg15446043, and *FOXO4* cg12453500 were related to both Fe and Ni. To illustrate the epigenome-wide association between $PM_{2.5}$ species and methylation, we plotted the -$\log_{10}$(*P*-value) from the mixed-effects models for all the 484,613 CpG sites in Fig. 2. The horizontal line represents the Bonferroni-corrected significance threshold. Subset analyses among nonsmokers and smokers showed consistent associations between PM species and methylation within smoking subgroups. For example, the coefficients for Fe at our most significant CpG (cg10911913) were 0.122 overall (Table S1 in the Supplemental Document), 0.131 among smokers and 0.086 among nonsmokers, and were statistically significant at a genome-wide level for both smoking and non-smoking subgroups. We observed the same trend for Ni and V.

Results of the bioinformatics analysis are shown in Fig. 3. We conducted KEGG pathway enrichment analysis

Table 3. Genome-wide associations of one-year moving averages of $PM_{2.5}$ species with DNA methylation: CpGs with Bonferroni-corrected statistical significance (*P*-value $< 9.4\times 10^{-9}$).

Species	Chr[a]	Gene	CpG	Location	Estimate[b]	Coef[c]	SE[d]	*P*-value
Fe	1	*DHRS3*	ch.1.471396R	Body	0.053	0.089	0.015	2.37E-09
	1	*CASZ1*	cg16238819	5′UTR	0.076	0.126	0.021	6.94E-09
	6	*SLC39A7;RXRB*	cg24319547	5′UTR;TSS1500	0.031	0.052	0.009	2.20E-09
	6	*KIAA1949*	cg08052856	5′UTR	0.072	0.120	0.020	2.58E-09
	6	*CDC5L*	cg22877480	TSS200	0.068	0.113	0.019	8.30E-09
	8	*FBXO25*	ch.8.7249R	Body	0.062	0.103	0.017	4.40E-09
	10	*CUEDC2*	cg06024834	TSS200	0.039	0.065	0.011	1.98E-09
	11	*RPS6KB2*	cg26457823	Body	0.035	0.059	0.010	3.82E-09
	11	*RPS25;TRAPPC4*	cg05223675	Body;TSS1500	0.077	0.128	0.021	6.97E-09
	14	*SMEK1*	cg01561368	TSS1500	0.024	0.040	0.007	6.20E-09
	15	*ONECUT1*	cg15446043	TSS200	0.146	0.243	0.039	1.37E-09
	15	*TPM1*	cg12302829	Body	0.039	0.065	0.011	2.72E-09
	17	*SLFN11*	cg10911913	TSS200	0.073	0.122	0.018	1.84E-10
	17	*CYTH1*	cg26525457	Body	−0.076	−0.126	0.020	1.26E-09
	18	*RNF138*	cg16078886	Body	0.065	0.108	0.018	2.55E-09
	19	*MYPOP*	cg03766703	3′UTR	−0.119	−0.198	0.033	6.91E-09
	19	*REEP6;PCSK4*	cg13970591	1stExon;TSS1500	0.222	0.370	0.063	8.10E-09
	21	*C21orf57*	cg10434947	5′UTR	0.043	0.072	0.012	4.32E-09
	22	*CBY1;LOC646851*	cg11003278	5′UTR;TSS1500	0.050	0.084	0.013	4.71E-10
	X	*FOXO4*	cg12453500	TSS1500	0.059	0.098	0.014	2.89E-11
Ni	1	*CASZ1*	cg16238819	5′UTR	0.032	1.062	0.179	8.25E-09
	10	*CUEDC2*	cg06024834	TSS200	0.017	0.560	0.089	1.10E-09
	15	*ONECUT1*	cg15446043	TSS200	0.059	1.950	0.330	8.31E-09
	16	*WDR90*	cg02734472	TSS200	0.035	1.168	0.194	4.61E-09
	17	*SLFN11*	cg10911913	TSS200	0.031	1.046	0.156	7.56E-11
	18	*RNF138*	cg16078886	Body	0.029	0.973	0.148	1.91E-10
	19	*SEC1;DBP*	cg13402773	TSS1500;TSS200	0.056	1.850	0.310	6.22E-09
	X	*FOXO4*	cg12453500	TSS1500	0.022	0.739	0.120	2.10E-09
V	17	*SLFN11*	cg10911913	TSS200	0.044	1.103	0.174	6.81E-10

[a]Chromosome.
[b]Estimate of change in methylation level per IQR increase in the species concentration.
[c]Regression coefficient in the mixed-effects models, with adjustment for $PM_{2.5}$ mass, Houseman estimated cell proportions ($CD4^+$ T lymphocytes, $CD8^+$ T lymphocytes, natural killer cells, B cells, monocytes), age, BMI, cigarette pack years, smoking status, alcohol consumption, education, year, season, and batch effects (plate, position of chip on plate, row and column position on chip).
[d]Standard error for regression coefficient.



Figure 2. Genome-wide associations of one-year moving averages of $PM_{2.5}$ species with DNA methylation, displayed by Manhattan plots. The horizontal line represents the significance threshold with Bonferroni correction (*P*-value $< 9.4 \times 10^{-9}$). We originally found 22 significant CpGs for Fe, 9 for Ni, and 1 for V, as shown in Figure 2. After excluding CpGs that were non-unimodal in the dip test, were cross-reactive, or contained SNPs in the last 10 bases, 20 CpGs remained significantly associated with Fe, 8 with Ni, and 1 with V.

for gene targets of CpGs with FDR <1% and <5%, respectively. Pathways whose enrichment FDRs were less than 5% were considered significant. In the analysis for gene targets of CpGs with FDR <1%, we identified 9 significant KEGG pathways. Three of them (i.e., "pathways in cancer," "non-small cell lung cancer," and "glioma") belong to the "Cancers" network. Results for gene targets of CpGs with FDR <5% suggest that those genes may also be linked to other cancers (e.g., myeloid leukemia) and pathways in the immune system. Other identified pathways include the diabetes and mitogen-activated protein kinases (MAPK) signaling pathways; the latter is involved in cellular functions including cell proliferation, differentiation, and migration.

Discussion

In this study, we investigated the relationship between $PM_{2.5}$ species and epigenome-wide DNA methylation in a large longitudinal cohort of elderly men. To the best of our knowledge, this is the first study that examined epigenome-wide associations between ambient $PM_{2.5}$ components and DNA methylation. We adjusted for multiple tests for the 11 PM components and 484,613 CpGs analyzed. We observed that 20 CpGs met the conservative Bonferroni-corrected level of significance (*P*-value $< 9.4 \times 10^{-9}$) in the analysis for Fe, 8 for Ni, and 1 for V. Fe mainly originates from crustal sources and the iron and steel industry; Ni and V are by-products of oil combustion, whereas



Figure 3. KEGG pathway enrichment analysis for (A) gene targets of CpGs with FDR < 1%, and (B) gene targets of CpGs with FDR < 5%. Pathways whose enrichment FDR was less than 5% were considered significant and are indicated by colored chunks.

Ni is also associated with the iron and steel industry.[40] The strong correlations between these species are shown in Table 2; the correlation coefficients were 0.93 for Fe-Ni, 0.86 for Fe-V, and 0.96 for Ni-V. In addition, we found that all 3 metals were associated with methylation levels at cg10911913 within the *SLFN11* gene. Our bioinformatics analysis suggests that gene targets of CpGs with FDR < 1% may be relevant to pathways including cancers, signal transduction, and cell growth and death.

To date, only a limited number of research studies have examined the association of ambient particles with DNA methylation. Most of them used surrogates for global DNA methylation, such as methylation at repetitive elements, or focused on methylation in candidate genes. The main shortcomings of these studies are: 1) methylation levels measured by surrogates do not consider the location in the genome and the patterns of methylation at surrogates may not well represent the patterns of gene-specific methylation, and 2) the use of candidate genes can only reveal methylation changes in particular genes selected *a priori* and may miss the genes that exhibit the most profound effects.

One previous study regarding PM exposure and DNA methylation reported that exposures to 90-day moving averages of black carbon and sulfate particles concentrations were associated with hypomethylation of the long interspersed nucleotide element-1 (LINE-1) and the short interspersed nucleotide element *Alu*.[6] Baccarelli et al.[41] found decreased LINE-1 methylation in association with 7-day moving average black carbon and $PM_{2.5}$ levels, whereas no association was observed for *Alu* methylation. A controlled human exposure experiment observed hypomethylation of *Alu* and the *toll-like receptor 4 (TLR4)* gene among 15 healthy adults exposed to fine concentrated ambient particles (CAPs) for 130 min.[13] Recently, a longitudinal study reported a relationship between decreased methylation at 3 CpG loci located in the *nitric oxide synthase 2, inducible (NOS2A)* gene and short-term exposures to organic carbon, elemental carbon, NO_3^-, and NH_4^+ among 28 patients with chronic obstructive pulmonary disease.[7]

A more recent epigenome-wide study by Panni et al. on short- and mid-term $PM_{2.5}$ exposure among 3 independent cohorts—the US NAS cohort and the German KORA F3 and F4 cohorts—identified 12 CpG sites at a Bonferroni adjusted

significance level.[20] Methylation at 9 out of these 12 CpGs showed increases related to PM exposure. This trend was consistent with our study: we found that 27 out of the 29 associations across all species displayed increased methylation levels. However, our study did not identify the same CpGs. A possible explanation for this difference is that the previous study evaluated short- or medium-term effects of PM, whereas we focused on long-term exposures. In addition, the results of the previous study were from a meta-analysis of 3 cohorts, including our study population. The 2 German cohorts had a much larger sample size than the NAS cohort (2299 vs. 657 participants), and the Panni et al. study reported heterogeneity at some CpG sites across the cohorts. Hence, their findings were more likely to reveal methylation changes within the German population and cannot be generalized to the US NAS population.

The most consistent finding of our study was the significant increase in methylation at *SLFN11* cg10911913 that was associated with one-year moving average concentrations of the $PM_{2.5}$ species Fe, Ni, and V. Schlafen Family Member *(SLFN)* genes encode a family of proteins that have been implicated in the regulation of cell growth and T cell development.[42,43] *SLFN11*, a member of the *SLFN* family, potently and specifically abrogates the production of retroviruses, e.g., human immunodeficiency virus (HIV-1).[33] Using cells with endogenously high and low *SLFN11* expression and siRNA-mediated silencing, Zoppoli et al.[44] found that *SLFN11* causally determines cell death and cell cycle arrest in response to DNA-damaging agents in cancer cells from different tissues of origin. *SLFN11* cg10911913 is located in a region whose function is promoter associated. Our results showed that methylation at this CpG site was positively associated with one-year moving averages of the $PM_{2.5}$ species Fe, Ni, and V. Our analysis suggests that exposure to ambient particles, especially those from oil combustion, was associated with hypermethylation of CpGs within *SLFN11*, which might lead to its downregulation. This hypothesis is consistent with recent literature on air-pollution-induced activation of viral sequences integrated in the human genome. For example, a study of truck drivers and office workers in Beijing, China showed a significant association of exposure to elemental carbon—taken as a marker of traffic particle exposure—with altered expression of viral microRNAs, possibly reflecting reactivation of latent retroviral sequences integrated in the host DNA.[45]

We also observed that methylation at *one cut homeobox 1 (ONECUT1)* cg15446043 was positively associated with Fe and Ni exposures. ONECUT1 can inhibit hepatitis B virus gene expression and DNA replication via both transcriptional and post-transcriptional mechanisms.[46]

In addition, we found that long-term exposure to Fe was linked to methylation in other genes expressed in immune cells, including *cytohesin 1 (CYTH1)* and *ribosomal protein S6 kinase (RPS6KB2). CYTH1* regulates adhesion and transendothelial migration of monocytes, T lymphocytes, and dentritic cells; it is also a key regulator of neutrophil adhesion to endothelial cells and to components of the extracellular matrix.[47,48] *RPS6KB2* plays an important role in neutrophilic differentiation and neutrophilic proliferation in human promyelocytic leukemia (HL-60) cells. Hence, methylation changes at pertinent CpG sites within these 2 genes may affect immune function.

Interestingly, we found that Fe was associated with methylation levels of CpGs in genes that regulate the cell cycle, such as *CDC5L* cg22877480 and *RPS6KB2* cg26457823.[49,50] *Cell division cycle 5-like (CDC5L)* is a crucial regulator of cell cycle G2/M progression and a component of pre-mRNA processing; hence, previous research has suggested it as a target for cancer therapy[50,51] RPS6KB2 is also highly active in the G2 and M phases.[49]

The DAVID bioinformatics analysis also indicated that Fe and Ni were associated with type II diabetes mellitus, insulin signaling pathways, and the MAPK pathway. This is consistent with previous work. For instance, long-term $PM_{2.5}$ exposure resulted in systemic inflammation, increased visceral adiposity, and insulin resistance among mice on a high fat diet.[52] In mice consuming normal diets, PM exposure was a risk factor for the development of type II diabetes, as it induced insulin resistance and impaired glucose tolerance.[53] Similarly, in humans, long-term exposure to particulate matter was linked to increased risk of type II diabetes in a large cohort study.[54] We have also reported that particulate air pollution was associated with changes in methylation in the MAPK pathway within the NAS cohort, but were unable to analyze metal components as in this study.[55]

Earlier, we found that particles from oil combustion were associated with markers of inflammation and endothelial dysfunction, and with increased blood pressure in the NAS population.[56,57]

The current study may indicate potential mechanisms of PM-related health effects on an epigenetic level: specifically, a proinflammatory stimulus such as air pollution may modify methylation in genes relevant to immune responses, inducing inflammation and endothelial dysfunction, thereby influencing cardiovascular health. Our study has several strengths. First, this is the first study that examined the effects of ambient particulate matter species on epigenome-wide DNA methylation. Previous studies often estimated DNA methylation associated with ambient PM using surrogates for global methylation or gene-specific methylation at candidate genes. An epigenome-wide study like ours offers a novel perspective on such an association, and implicates potential biologic mechanisms on an epigenetic level. Second, we used a stringent Bonferroni correction threshold to adjust the level of significance, which accounted both for the 11 $PM_{2.5}$ species and for the number of CpG sites analyzed. Although we observed a small number of significant CpGs due to the strict correction, the probability of false positives is expected to remain appropriate. Third, the study population was geographically stable and well followed since enrollment in 1,963.

On the other hand, this study has several notable limitations. One major limitation is that we were unable to perform a replication analysis due to a lack of appropriate comparison cohorts. The German KORA cohorts had $PM_{2.5}$ mass data but did not have measurements on PM species. Since the chemical composition of air pollutants in Germany may differ from that in the US, we were not able to impute species data from $PM_{2.5}$ measurements. We do not have information on gene expression, and thus we cannot confirm the expected functional

implications of the identified methylation changes. Further research on gene expression in relation to PM exposures is needed. Additionally, we are unable to rule out residual confounding. The bioinformatics analysis of pathways used annotations of methylation sites within genes and thus genes that include more sites on the 450K array, such as those involved in cancer, are more likely to arise in pathway results.[58] Last but not least, there exists a generalizability issue because our study population consisted almost entirely of elderly Caucasian men. One should consider factors like age, sex, and race when generalizing our findings to other populations.

In summary, we examined the association of long-term exposure to $PM_{2.5}$ species with differential DNA methylation in a 450K epigenome-wide association study using a large longitudinal cohort. Our study added evidence on the association of ambient particulate exposure with epigenome-wide DNA methylation.[20] It also suggests differing relative toxicities of PM species, which is especially important given the US. National Research Council's recommendation to consider determining the differential risk of $PM_{2.5}$ species a research priority.[59] Furthermore, loci identified in the study provide a possible link between $PM_{2.5}$ and its identified health effects. Specifically, the current study suggests that particles from oil combustion are associated with methylation changes in genes relevant to immune responses. Further studies are needed to explore the mechanisms behind the relationship between ambient particles and DNA methylation.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

This work was supported by the National Institute of Environmental Health Sciences ES000002, ES021733 and ES015172–01 and the US. Environmental Protection Agency RD-83587201. A.J. was supported by the National Institute of Environmental Health Sciences K99 ES023450. D.S. was supported by a VA Research Career Scientist Award. The VA Normative Aging Study is supported by the Cooperative Studies Program/Epidemiology Research and Information Centers of the US. Department of Veterans Affairs and is a component of the Massachusetts Veterans Epidemiology Research and Information Center, Boston, Massachusetts.

Disclaimers

The contents are solely the responsibility of the grantee and do not necessarily represent the official views of the funders. Further, the funders do not endorse the purchase of any commercial products or services mentioned in the publication.

ORCID

Allan C. Just http://orcid.org/0000-0003-4312-5957
Andrea A. Baccarelli http://orcid.org/0000-0002-3436-0640

References

1. Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A. The contribution of outdoor air pollution sources to premature mortality on a global scale. Nature 2015; 525:367-71; PMID:26381985; http://dx.doi.org/10.1038/nature15371
2. Zanobetti A, Franklin M, Koutrakis P, Schwartz J. Fine particulate air pollution and its components in association with cause-specific emergency admissions. Environ Health 2009; 8:58; PMID:20025755; http://dx.doi.org/10.1186/1476-069X-8-58
3. Dai L, Zanobetti A, Koutrakis P, Schwartz JD. Associations of fine particulate matter species with mortality in the United States: a multicity time-series analysis. Environ Health Perspect 2014; 122:837-42; PMID:24800826; http://dx.doi.org/10.1289/ehp.1307568
4. Krall JR, Anderson GB, Dominici F, Bell ML, Peng RD. Short-term exposure to particulate matter constituents and mortality in a national study of U.S. urban communities. Environ Health Perspect 2013; 121:1148-53; PMID:23912641; http://dx.doi.org/10.1289/ehp.1206185
5. Peng R, Bell M, Geyh A, McDermott A, Zeger S, Samet J, Dominici F. Emergency admissions for cardiovascular and respiratory diseases and the chemical composition of fine particle air pollution. Environ Health Perspect 2009; 117:957-63; PMID:19590690; http://dx.doi.org/10.1289/ehp.0800185
6. Madrigano J, Baccarelli A, Mittleman MA, Wright RO, Sparrow D, Vokonas PS, Tarantini L, Schwartz J. Prolonged exposure to particulate pollution, genes associated with glutathione pathways, and DNA methylation in a cohort of older men. Environ Health Perspect 2011; 119:977-82; PMID:21385671; http://dx.doi.org/10.1289/ehp.1002773
7. Chen R, Qiao L, Li H, Zhao Y, Zhang Y, Xu W, Wang C, Wang H, Zhao Z, Xu X, et al. Fine Particulate Matter Constituents, Nitric Oxide Synthase DNA Methylation and Exhaled Nitric Oxide. Environ Sci Technol 2015; 49:11859-65; PMID:26372312; http://dx.doi.org/10.1021/acs.est.5b02527
8. Baccarelli A, Bollati V. Epigenetics and environmental chemicals. Curr Opin Pediatr 2009; 21:243-51; PMID:19663042; http://dx.doi.org/10.1097/MOP.0b013e32832925cc
9. Belinsky SA, Palmisano WA, Gilliland FD, Crooks LA, Divine KK, Winters SA, Grimes MJ, Harms HJ, Tellez CS, Smith TM, et al. Aberrant promoter methylation in bronchial epithelium and sputum from current and former smokers. Cancer Res 2002; 62:2370-7; PMID:11956099
10. Chen H, Li S, Liu J, Diwan BA, Barrett JC, Waalkes MP. Chronic inorganic arsenic exposure induces hepatic global and individual gene hypomethylation: implications for arsenic hepatocarcinogenesis. Carcinogenesis 2004; 25:1779-86; PMID:15073043; http://dx.doi.org/10.1093/carcin/bgh161
11. Baccarelli A, Tarantini L, Wright RO, Bollati V, Litonjua AA, Zanobetti A, Sparrow D, Vokonas PS, Schwartz J. Repetitive element DNA methylation and circulating endothelial and inflammation markers in the VA normative aging study. Epigenetics 2010; 5:222-8; PMID:20305373; http://dx.doi.org/10.4161/epi.5.3.11377
12. Castro R, Rivera I, Struys EA, Jansen EE, Ravasco P, Camilo ME, Blom HJ, Jakobs C, Tavares de Almeida I. Increased homocysteine and S-adenosylhomocysteine concentrations and DNA hypomethylation in vascular disease. Clin Chem 2003; 49:1292-6; PMID:12881445; http://dx.doi.org/10.1373/49.8.1292
13. Bellavia A, Urch B, Speck M, Brook RD, Scott JA, Albetti B, Behbod B, North M, Valeri L, Bertazzi PA, et al. DNA hypomethylation, ambient particulate matter, and increased blood pressure: findings from controlled human exposure experiments. J Am Heart Assoc 2013; 2:e000212; PMID:23782920; http://dx.doi.org/10.1161/JAHA.113.000212
14. Madrigano J, Baccarelli A, Mittleman MA, Sparrow D, Spiro A, 3rd, Vokonas PS, Cantone L, Kubzansky L, Schwartz J. Air pollution and DNA methylation: interaction by psychological factors in the VA Normative Aging Study. Am J Epidemiol 2012; 176:224-32; PMID:22798479; http://dx.doi.org/10.1093/aje/kwr523
15. Fan T, Fang SC, Cavallari JM, Barnett IJ, Wang Z, Su L, Byun HM, Lin X, Baccarelli AA, Christiani DC. Heart rate variability and DNA methylation levels are altered after short-term metal fume exposure among occupational welders: a repeated-measures panel study. BMC Public Health 2014; 14:1279; PMID:25512264; http://dx.doi.org/10.1186/1471-2458-14-1279
16. Kile ML, Fang S, Baccarelli AA, Tarantini L, Cavallari J, Christiani DC. A panel study of occupational exposure to fine particulate matter and changes in DNA methylation over a single workday and years

worked in boilermaker welders. Environ Health 2013; 12:47; PMID:23758843; http://dx.doi.org/10.1186/1476-069X-12-47
17. Joubert BR, Haberg SE, Nilsen RM, Wang X, Vollset SE, Murphy SK, Huang Z, Hoyo C, Midttun O, Cupul-Uicab LA, et al. 450K epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy. Environ Health Perspect 2012; 120:1425-31; PMID:22851337; http://dx.doi.org/10.1289/ehp.1205412
18. Joubert BR, Felix JF, Yousefi P, Bakulski KM, Just AC, Breton C, Reese SE, Markunas CA, Richmond RC, Xu CJ, et al. DNA Methylation in Newborns and Maternal Smoking in Pregnancy: Genome-wide Consortium Meta-analysis. Am J Hum Genet 2016; 98:680-96; PMID:27040690; http://dx.doi.org/10.1016/j.ajhg.2016.02.019
19. Zeilinger S, Kuhnel B, Klopp N, Baurecht H, Kleinschmidt A, Gieger C, Weidinger S, Lattka E, Adamski J, Peters A, et al. Tobacco smoking leads to extensive genome-wide changes in DNA methylation. PLoS One 2013; 8:e63812; PMID:23691101; http://dx.doi.org/10.1371/journal.pone.0063812
20. Panni T, Mehta AJ, Schwartz JD, Baccarelli AA, Just AC, Wolf K, Wahl S, Cyrys J, Kunze S, Strauch K, et al. A Genome-Wide Analysis of DNA Methylation and Fine Particulate Matter Air Pollution in Three Study Populations: KORA F3, KORA F4, and the Normative Aging Study. Environ Health Perspect 2016; 124(7):983-90; PMID:26731791; http://dx.doi.org/10.1289/ehp.1509966
21. Bell B, Rose C, Damon A. The Normative Aging Study: an interdisciplinary and longitudinal study of health and aging. Aging Hum Dev 1972; 3:4-17; http://dx.doi.org/10.2190/GGVP-XLB5-PC3N-EF0G
22. Bibikova M, Barnes B, Tsan C, Ho V, Klotzle B, Le JM, Delano D, Zhang L, Schroth GP, Gunderson KL, et al. High density DNA methylation array with single CpG site resolution. Genomics 2011; 98:288-95; PMID:21839163; http://dx.doi.org/10.1016/j.ygeno.2011.07.007
23. Triche TJ, Jr., Weisenberger DJ, Van Den Berg D, Laird PW, Siegmund KD. Low-level processing of Illumina Infinium DNA Methylation BeadArrays. Nucleic Acids Res 2013; 41:e90; PMID:23476028; http://dx.doi.org/10.1093/nar/gkt090
24. Davis S, Du P, Bilke S, Triche T, Jr., Bootwalla M. methylumi: Handle Illumina methylation data. R package version 2.17.0., 2015
25. Teschendorff AE, Jones A, Fiegl H, Sargent A, Zhuang JJ, Kitchener HC, Widschwendter M. Epigenetic variability in cells of normal cytology is associated with the risk of future morphological transformation. Genome Med 2012; 4:24; PMID:22453031; http://dx.doi.org/10.1186/gm323
26. Zhu ZZ, Hou L, Bollati V, Tarantini L, Marinelli B, Cantone L, Yang AS, Vokonas P, Lissowska J, Fustinoni S, et al. Predictors of global methylation levels in blood DNA of healthy subjects: a combined analysis. Int J Epidemiol 2012; 41:126-39; PMID:20846947; http://dx.doi.org/10.1093/ije/dyq154
27. Mostofsky E, Schwartz J, Coull BA, Koutrakis P, Wellenius GA, Suh HH, Gold DR, Mittleman MA. Modeling the association between particle constituents of air pollution and health outcomes. Am J Epidemiol 2012; 176:317-26; PMID:22850792; http://dx.doi.org/10.1093/aje/kws018
28. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, Wiencke JK, Kelsey KT. DNA methylation arrays as surrogate measures of cell mixture distribution. BMC Bioinformatics 2012; 13:86; PMID:22568884; http://dx.doi.org/10.1186/1471-2105-13-86
29. Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, Irizarry RA. Minfi: a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. Bioinformatics (Oxford, England) 2014; 30:1363-9; PMID:24478339; http://dx.doi.org/10.1093/bioinformatics/btu049
30. Reinius LE, Acevedo N, Joerink M, Pershagen G, Dahlen SE, Greco D, Soderhall C, Scheynius A, Kere J. Differential DNA methylation in purified human blood cells: implications for cell lineage and studies on disease susceptibility. PLoS One 2012; 7:e41361; PMID:22848472; http://dx.doi.org/10.1371/journal.pone.0041361
31. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. Nat Protoc 2009; 4:44-57; PMID:19131956; http://dx.doi.org/10.1038/nprot.2008.211
32. Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res 2009; 37:1-13; PMID:19033363; http://dx.doi.org/10.1093/nar/gkn923
33. Li M, Kao E, Gao X, Sandig H, Limmer K, Pavon-Eternod M, Jones TE, Landry S, Pan T, Weitzman MD, et al. Codon-usage-based inhibition of HIV protein synthesis by human schlafen 11. Nature 2012; 491:125-8; PMID:23000900; http://dx.doi.org/10.1038/nature11433
34. Hartigan JA, Hartigan PM. The Dip Test of Unimodality. Ann Statist 1985; 13:70-84; http://dx.doi.org/10.1214/aos/1176346577
35. Chen YA, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW, Gallinger S, Hudson TJ, Weksberg R. Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium HumanMethylation450 microarray. Epigenetics 2013; 8:203-9; PMID:23314698; http://dx.doi.org/10.4161/epi.23470
36. Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, Handsaker RE, Kang HM, Marth GT, McVean GA. An integrated map of genetic variation from 1,092 human genomes. Nature 2012; 491:56-65; PMID:23128226; http://dx.doi.org/10.1038/nature11632
37. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/. 2014
38. Kanehisa M, Goto S, Kawashima S, Nakaya A. The KEGG databases at GenomeNet. Nucleic Acids Res 2002; 30:42-6; PMID:11752249; http://dx.doi.org/10.1093/nar/30.1.42
39. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. J Royal Statist Soc Series B (Methodological) 1995; 57:289-300
40. Geiger A, Cooper J. Overview of Airborne Metals Regulations, Exposure Limits, Health Effects, and Contemporary Research. 2010
41. Baccarelli A, Wright RO, Bollati V, Tarantini L, Litonjua AA, Suh HH, Zanobetti A, Sparrow D, Vokonas PS, Schwartz J. Rapid DNA methylation changes after exposure to traffic particles. Am J Respir Crit Care Med 2009; 179:572-8; PMID:19136372; http://dx.doi.org/10.1164/rccm.200807-1097OC
42. Geserick P, Kaiser F, Klemm U, Kaufmann SH, Zerrahn J. Modulation of T cell development and activation by novel members of the Schlafen (slfn) gene family harbouring an RNA helicase-like motif. Inter Immunol 2004; 16:1535-48; PMID:15351786; http://dx.doi.org/10.1093/intimm/dxh155
43. Schwarz DA, Katayama CD, Hedrick SM. Schlafen, a new family of growth regulatory genes that affect thymocyte development. Immunity 1998; 9:657-68; PMID:9846487; http://dx.doi.org/10.1016/S1074-7613(00)80663-9
44. Zoppoli G, Regairaz M, Leo E, Reinhold WC, Varma S, Ballestrero A, Doroshow JH, Pommier Y. Putative DNA/RNA helicase Schlafen-11 (SLFN11) sensitizes cancer cells to DNA-damaging agents. Proc Natl Acad Sci 2012; 109:15030-5; PMID:22927417; http://dx.doi.org/10.1073/pnas.1205943109
45. Hou L, Barupal J, Zhang W, Zheng Y, Liu L, Zhang X, Dou C, McCracken JP, Diaz A, Motta V, et al. Particulate Air Pollution Exposure and Expression of Viral and Human MicroRNAs in Blood: The Beijing Truck Driver Air Pollution Study. Environ Health Perspect 2015; 124(3):344-50; PMID:26068961; http://dx.doi.org/10.1289/ehp.1408519
46. Hao R, He J, Liu X, Gao G, Liu D, Cui L, Yu G, Yu W, Chen Y, Guo D. Inhibition of hepatitis B virus gene expression and replication by hepatocyte nuclear factor 6. J Virol 2015; 89:4345-55; PMID:25653429; http://dx.doi.org/10.1128/JVI.03094-14
47. El azreq MA, Bourgoin SG. Cytohesin-1 regulates human blood neutrophil adhesion to endothelial cells through beta2 integrin activation. Mol Immunol 2011; 48:1408-16; PMID:21511340; http://dx.doi.org/10.1016/j.molimm.2011.03.018
48. Kolanus W. Guanine nucleotide exchange factors of the cytohesin family and their roles in signal transduction. Immunol Rev 2007; 218:102-13; PMID:17624947; http://dx.doi.org/10.1111/j.1600-065X.2007.00542.x
49. Boyer D, Quintanilla R, Lee-Fruman KK. Regulation of catalytic activity of S6 kinase 2 during cell cycle. Mol Cell Biochem 2008; 307:59-64; PMID:17786541; http://dx.doi.org/10.1007/s11010-007-9584-5

50. Graub R, Lancero H, Pedersen A, Chu M, Padmanabhan K, Xu XQ, Spitz P, Chalkley R, Burlingame AL, Stokoe D, et al. Cell cycle-dependent phosphorylation of human CDC5 regulates RNA processing. Cell cycle (Georgetown, Tex) 2008; 7:1795-803; PMID:18583928; http://dx.doi.org/10.4161/cc.7.12.6017
51. Mu R, Wang YB, Wu M, Yang Y, Song W, Li T, Zhang WN, Tan B, Li AL, Wang N, et al. Depletion of pre-mRNA splicing factor Cdc5L inhibits mitotic progression and triggers mitotic catastrophe. Cell death & disease 2014; 5:e1151; PMID:24675469; http://dx.doi.org/10.1038/cddis.2014.117
52. Sun Q, Yue P, Deiuliis JA, Lumeng CN, Kampfrath T, Mikolaj MB, Cai Y, Ostrowski MC, Lu B, Parthasarathy S, et al. Ambient air pollution exaggerates adipose inflammation and insulin resistance in a mouse model of diet-induced obesity. Circulation 2009; 119:538-46; PMID:19153269; http://dx.doi.org/10.1161/CIRCULATIONAHA.108.799015
53. Xu X, Liu C, Xu Z, Tzan K, Zhong M, Wang A, Lippmann M, Chen LC, Rajagopalan S, Sun Q. Long-term exposure to ambient fine particulate pollution induces insulin resistance and mitochondrial alteration in adipose tissue. Toxicol Sci 2011; 124:88-98; PMID:21873646; http://dx.doi.org/10.1093/toxsci/kfr211
54. Weinmayr G, Hennig F, Fuks K, Nonnemacher M, Jakobs H, Mohlenkamp S, Erbel R, Jockel KH, Hoffmann B, Moebus S. Long-term exposure to fine particulate matter and incidence of type 2 diabetes mellitus in a cohort study: effects of total and traffic-specific air pollution. Environ Health 2015; 14:53; PMID:26087770; http://dx.doi.org/10.1186/s12940-015-0031-x
55. Carmona JJ, Sofer T, Hutchinson J, Cantone L, Coull B, Maity A, Vokonas P, Lin X, Schwartz J, Baccarelli AA. Short-term airborne particulate matter exposure alters the epigenetic landscape of human genes associated with the mitogen-activated protein kinase network: a cross-sectional study. Environ Health 2014; 13:94; PMID:25395096; http://dx.doi.org/10.1186/1476-069X-13-94
56. Dai L, Bind MA, Koutrakis P, Coull BA, Sparrow D, Vokonas PS, Schwartz JD. Fine particles, genetic pathways, and markers of inflammation and endothelial dysfunction: Analysis on particulate species and sources. J Expo Sci Environ Epidemiol 2016; 26(4):415-21; PMID:26732377; http://dx.doi.org/10.1038/jes.2015.83
57. Dai L, Koutrakis P, Coull BA, Sparrow D, Vokonas PS, Schwartz JD. Use of the Adaptive LASSO Method to Identify PM Components Associated with Blood Pressure in Elderly Men: The Veterans Affairs Normative Aging Study. Environ Health Perspect 2015; 124(1):120-5; PMID:26090776; http://dx.doi.org/10.1289/ehp.1409021
58. Harper KN, Peters BA, Gamble MV. Batch effects and pathway analysis: 2 potential perils in cancer studies involving DNA methylation array analysis. Cancer Epidemiol Biomarkers Prev 2013; 22:1052-60; http://dx.doi.org/10.1158/1055-9965.EPI-13-0114
59. National Reserch Council. Research Priorities for Airborne Particulate Matter: IV. Continuing Research Progress Washington, DC: National Research Council 2004